



FREE NEJM E-TOC

HOME

SUBSCRIBE

CURRENT ISSUE

PAST ISSUES

COLLECTIONS

Keyword, citation, or authc

UNIVERSITY OF SOUTH ALABAMA | [Get NEJM's E-Mail Table of Contents - FREE](#) | [Sign In as Individual](#) | [Contact Administrator](#)

ORIGINAL ARTICLE

Beckett et al. 358 (18): 1887, Table 2 May 1, 2008

▶ [Return to Article](#)

▶ [Add to Personal Archive](#)

▶ [PowerPoint Help](#)

[Get PowerPoint Slide ▶](#)

Table 2. Main Fatal and Nonfatal End Points in the Intention-to-Treat Population.

End Point	Rate per 1000 Patient-Yr (No. of Events)		Unadjusted Hazard Ratio (95% CI)	P Value
	Active	Placebo		
	<i>no. (%)</i>			
Stroke				
Fatal or nonfatal	12.4 (51)	17.7 (69)	0.70 (0.49–1.01)	0.06
Death from stroke	6.5 (27)	10.7 (42)	0.61 (0.38–0.99)	0.046
Death				
From any cause	47.2 (196)	59.6 (235)	0.79 (0.65–0.95)	0.02
From noncardiovascular or unknown causes	23.4 (97)	28.9 (114)	0.81 (0.62–1.06)	0.12
From cardiovascular cause	23.9 (99)	30.7 (121)	0.77 (0.60–1.01)	0.06
From cardiac cause*	6.0 (25)	8.4 (33)	0.71 (0.42–1.19)	0.19
From heart failure	1.5 (6)	3.0 (12)	0.48 (0.18–1.28)	0.14
Fatal or nonfatal				
Any myocardial infarction	2.2 (9)	3.1 (12)	0.72 (0.30–1.70)	0.45
Any heart failure	5.3 (22)	14.8 (57)	0.36 (0.22–0.58)	<0.001
Any cardiovascular event†	33.7 (138)	50.6 (193)	0.66 (0.53–0.82)	<0.001

* Death from cardiac causes was defined as fatal myocardial infarction, fatal heart failure, and sudden death.

† Any cardiovascular event was defined as death from cardiovascular causes or stroke, myocardial infarction, or heart failure.

Table 2. Main Fatal and Nonfatal End Points in the Intention-to-Treat Population.

[HOME](#) | [SUBSCRIBE](#) | [SEARCH](#) | [CURRENT ISSUE](#) | [PAST ISSUES](#) | [COLLECTIONS](#) | [PRIVACY](#) | [HELP](#) | [beta.nejm.org](#)

Comments and questions? Please [contact us](#).

The New England Journal of Medicine is owned, published, and [copyrighted](#) © 2008 [Massachusetts Medical Society](#). All rights reserved.